

## INHALTSVERZEICHNIS

- |                    |                                |   |
|--------------------|--------------------------------|---|
| Thyroid-Lit. 50986 | BROSE, M.S. et al.             | Multikinase inhibitors for the treatment of asymptomatic radioactive iodine-refractory differentiated thyroid cancer: global noninterventional study (RIFTOS MKI)         |
| Thyroid-Lit. 50987 | PALERMO, A. et al.             | Clinical use of Raman spectroscopy improves diagnostic accuracy for indeterminate thyroid nodules.  |
| Thyroid-Lit. 50988 | NOTO, B. et al.                | Predictors of anxiety and depression in differentiated thyroid cancer survivors: results of a cross-sectional study.  |
| Thyroid-Lit. 50989 | KEEFE, G. et al.               | Autoimmune thyroiditis and risk of malignancy in children with thyroid nodules.   |
| Thyroid-Lit. 50990 | GILL, S. et al.                | Evaluating the progression to hypothyroidism in preconception euthyroid thyroid peroxidase antibody-positive women.   |
| Thyroid-Lit. 50991 | ORSOLINI, F. et al.            | Thyroid function rather than thyroid antibodies affects pregnancy and perinatal outcomes: results of a prospective study.   |
| Thyroid-Lit. 50992 | DU PUY, R.S. et al.            | No effect of levothyroxine on hemoglobin in older age with subclinical hypothyroidism: pooled results from 2 randomized clinical trials.                                  |
| Thyroid-Lit. 50993 | KRIEGE, M. et al.              | Impact of direct laryngoscopy vs. videolaryngoscopy on signal quality of recurrent laryngeal nerve monitoring in thyroid surgery: a randomised parallel group trial.      |
| Thyroid-Lit. 50994 | DA SILVEIRA DUVAL, M.A. et al. | An undetectable postoperative calcitonin level is associated with long-term disease-free survival in medullary thyroid carcinoma: Results of a retrospective cohort study |

Thyroid-Lit. 50995	JERKOVICH, F. et al.	Active surveillance of small metastatic lymph nodes as an alternative to surgery in selected patients with low-risk papillary thyroid cancer: A retrospective cohort study
Thyroid-Lit. 50996	SAADAT, N. et al.	Treatment of post-radioactive iodine relapse of hyperthyroidism: comparison of long-term methimazole and radioactive iodine treatment
Thyroid-Lit. 50997	DUCHARME, M. et al.	A Novel Levothyroxine solution results in similar bioavailability whether taken 30 or just 15 minutes before a high-fat high-calorie meal